
LIFE SCIENCE AND HUMANITIES

Manuscript info:

Received November 4, 2018., Accepted November 17, 2018., Published November 30, 2018.

**IMPACT OF ANTHROPOGENIC DISTURBANCE ON
ANURANS HABITAT AND SPECIES RICHNESS IN
SILAGO, SOUTHERN LEYTE, PHILIPPINES**

WILBERT A. AUREO,

Forestry and Environmental Science Department,
College of Agriculture and Natural Resources,
Bohol Island State University,
Bilar Bohol. Email: waureo@vsu.edu.ph

MARLITO M. BANDE,

Institute of Tropical Ecology and Environmental Management,
College of Forestry and Environmental Science, Visayas State
University, ViscaBaybay, Leyte



<http://dx.doi.org/10.26739/2573-5616-2018-12-22>

Abstract: Studying the response of anurans to anthropogenic disturbance can be very useful in determining the potentiality of this species as bio-indicator. The study was conducted to assess and describe the habitat condition (i.e., microclimatic, microhabitat, and vegetation structures) and species richness of anurans as affected by the degree of disturbance due to anthropogenic activities in Silago, Southern Leyte, Philippines. Four (10 x 10 m) quadrats were established in each site using opportunistic approach in the collection of anurans within the plots per site. Likewise, measurement on the factors associated with habitat condition such as microclimate, habitat, and vegetation structure were undertaken. The collection was done both in daytime and nighttime. The results showed that microclimate, habitat, and vegetation structure differed significantly ($p < 0.01$) in each site. This implies that the three sites had different degrees of disturbance that altered the habitat condition of anurans. Moreover, anuran species richness significantly differed ($p < 0.05$) in primary forest and slash and burn cultivation but there was no significant difference between secondary and primary forest. Therefore, the results of this study may indicate that the degree of anthropogenic disturbance considerably ($p < 0.01$) alters anuran's habitat conditions which significantly ($p < 0.05$) affect its species richness.

Keywords: Tropical rainforest, ecological disturbance, anthropogenic activities, anurans, conservation

Recommended citation: WILBERT A. AUREO, MARLITO M. BANDE. IMPACT OF ANTHROPOGENIC DISTURBANCE ON ANURANS HABITAT AND SPECIES RICHNESS IN SILAGO, SOUTHERN LEYTE, PHILIPPINES. 11-12. American Journal of Research P. 198-208 (2018).

INTRODUCTION

Anurans or frogs are tailless amphibians whose life cycles include aquatic and terrestrial phases. Early in their development post-hatching, tadpoles survive by eating muck or small aquatic insects, until they finally metamorphose, and only then do they come onto land (Heying, 2004). They are highly sensitive and vulnerable to environmental changes and changes to habitat quality and are extremely sensitive to small changes in temperature and moisture because of their permeable skin and unshelled eggs (Cushman, 2006). Breeding is influenced by rainfall, humidity and temperature, and so any change or alteration in these abiotic factors leads to disruption of lifecycle. This is one reason why anurans, like other amphibians, are directly affected by any environmental stress in their terrestrial or aquatic habitats. In fact, studies have shown that most anurans located in polluted aquatic ecosystem exhibit malformation of organs and other body parts (Dey, 2010).

The combinations of natural and anthropogenic factors such as climate, geographic ranges and vegetation types (Cueto and Casenave, 1999) are also considered to have a significant influence on species distributions and diversity (Mallari, 2009). The pattern of increase in habitat heterogeneity on the structure and composition of vegetation becomes complex if the niche diversity and species diversity increases (Cramer and Willig, 2002; Tews et al., 2004; Gingold et al.,

2010; MacClain and Barry, 2010). Moreover, several species of forest anurans are correlated with the quantity and quality of woody debris, litter depth, and over-storey canopy closure (deMaynadier and Hunter, 1995).

The Philippines is one of the few countries in the world that is covered by rain forest (Schulte, 2002). It is also considered as one of the world's mega diverse countries that hosts a large share of endemic flora and fauna (Myer et al., 2000). However, due to the conversion of forests to marginal agriculture, commercial agriculture, and timber plantations, these forest resources are disappearing at an alarming rate (Margraf and Milan, 2004.). Filipino farmers who are poor and lack employment opportunities in the lowland migrate to the upland areas where they cut down secondary forest and practice kaingin farming (Kummer, 1992).

Silago forest is one of the remaining primary forests in the region. However due to rampant and uncontrolled hunting of wildlife and habitat destruction caused by slash-and-burn cultivation or kaingin contributed by residents both within and outside the communities, threat on biodiversity increases (Ceniza et al., 2011). In Barangays Imelda, Katipunan and Catmon, slash-and-burn cultivation is being practiced by the residents. This is their major source of livelihood because many of them do not own land in the alienable and disposable areas.

Thus, the current study examines the impact of ecological disturbance to habitat condition (i.e. rainfall,

temperature, relative humidity, vegetation structure, litter depth, and woody debris) and anurans species and population due to anthropogenic activity.

MATERIALS AND METHOD

Description of Study Sites

The study was conducted in Brgy. Katipunan, Silago Southern Leyte. The study site is part of the Mt. Nacolod Mountain Range and the last remaining primary forest on Leyte Island. The criteria in selecting the site is the degree of ecological disturbance following Miller (1982). Hence, the following sites were selected: primary forest, secondary forest, and slash and burn cultivation. (Fig 1).

Primary Forest

The area is part of the remaining primary forest of Mt Nacolod mountain range. The dominant vegetation cover in this area are mostly under the family of Dipterocarpacea specifically tanguile (*Shorea polysperma*, Blanco Merr) and red lauán (*Shorea negrosensis*, Foxw). Understorey is being dominated by Dipterocarpacea and Lauraceae species. Moreover, ground cover vegetation of the area is being dominated by ferns and tree wildlings. Remnants of cut trees were observed in the area.


Secondary Forest

The area is a secondary growth forest under early successional stage. This site was being cleared in year 2010. However, pioneer trees started to grow and dominate the area; thus, stabilizing the ecosystem. The vegetation that dominates the crown

cover is under Euphorbiaceae and Moraceae family specifically *Hinlaumong-puti* (*Mallotus paniculatos* Elm.), *Macaranga* (*Macaranga caudatifolia* Elm.) and *Ficus* sp. Moreover, grasses, ferns, and some wildlings of *Ficus* and *Macaranga* dominate the ground layer.

Slash and Burn Cultivation

The area is a newly cultivated slash-and-burn site. This is being managed and cultivated by an individual farmer. Tree stumps and partially burned logs were observed in the area. During the first and second collection, no vegetation was observed. However, on the third collection, the farmer started to plant cassava and coconut. Grasses and other herbaceous plants started to grow.



Note: (Site 1 - Primary forest; Site 2 - Secondary forest and Site 3 - slash and burn cultivation)

Figure 1. Location of the sampling plots that were randomly established within the three

selected study sites in Brgy. Katipunan, Silago, Southern Leyte

Sampling Method

Figure 1 presents the four (10 x 10 m) quadrats that were randomly

established in each site following the methods of Williams (2004). Anurans were collected at early morning 6:00-9:00 am and evening at 7:00-10:00 pm (Warguez et al, 2013). The Visual Encounter Survey was used to search high potential areas throughout the sampling sites. These are on the surface and under rocks, logs, trees, and other debris within each established plot.

Measurement of Anurans Morphometric

The following morphometric measurements were taken using a vernier caliper to the nearest centimeter to support identification of anurans to the species level following Alcalá and Brown (1998) and Alcalá (1996) as adopted by HARIBON Foundation guidelines for amphibians. Morphometric includes Snout-Vent Length (SVL), a measurement from the snout to the tip of the ventral part; TibiaLength (TBL), from the tibiato the tip of the phalanges (leg not straightened); Head Length (HdL), from the snout to the tympanum; Hind limb Length (HbL), entire length of the hind limb (straightened); Eye Diameter (ED), diameter of the entire eye; Tympanum Distance (TD), distance of tympanum from both ends; Snout Length (SL), from the snout to the eye; and Head Breadth, from one shoulder to the other.

Identification of Anurans

The morphometric of each captured frog were measured. This was done in order to support the identification of anurans. Likewise, photographs of each captured anurans were taken using a digital camera. Pre-identification was done

using the field guide of Diesmos and Alcalá (2011). Moreover, photos and morphometric measurements were sent to an expert to confirm preliminary identification. Lastly, final confirmation was done using IUCN List of Threatened Species available at (<http://www.iucnredlist.org/amphibians>, 2015).

Assessment of Vegetation Structure

Trees and other associated plants inside the plot were counted based on the following vegetation strata: a) Dominant, b) understorey vegetation, c) ground cover. Dominant stratum are the vegetation present in the uppermost stratum within a plot. Trees with diameter at breast height >15 cm and height >20 m were categorized as dominant tree. Understorey vegetation is the stratum next to the dominant layer which usually consists of shrubs and bushes. Trees with diameter at breast height <15 cm and height <20 m were categorized under understorey layer. The herbaceous layer is the vegetation present in the ground covering the soil such as grasses and ferns. Taxonomic identification of trees, associated ferns, grass, palms, and rattan was done at genus level.

Determination of Microhabitat Structure

Quantity and quality of Coarse Woody Debris (CWD) were determined by counting all CWD with small-end diameter (<10 cm) and greater than or equal to 10 cm that fell inside the quadrat. Each piece was rated on its degree of decomposition (Woodwall and Willams, 2005) ranging from 1

(sound, intact, no rot) to 5 (no structural integrity, soft, powdery). Also, litter depth in each quadrat was measured using a meter stick by randomly selecting three points of location per quadrat.

Measurement of Microclimatic Factors

Microclimatic factors measured were air temperature, relative humidity, and rainfall. Air temperature was taken at each quadrat per study site using a thermometer in an interval of 3 hours starting from 7:00 am to 4:00 pm. Relative humidity and rainfall were measured using a sling psychrometer and improvised rain gauge, respectively (with the same collection time and interval with the air temperature measurement).

Calculation of Species Richness

Calculations of species richness was done per plot and study site using the formula (Magurran, 1988):

$$S = \text{Total number of species}$$

Equation 1

Statistical Analysis

All data were tested for normality and heterogeneity using PROC Univariate of Statistical Analysis System version 9.1 (SAS, 2003). PROC GLM (general linear model) procedure was initially performed to check for effects of ecological disturbance on microclimatic factors (i.e., temperature, relative humidity, and rainfall), habitat structure (i.e., litter depth and quantity and quality of coarse woody debris), vegetation structure (i.e., herbaceous layer, understorey, and climax strata), and species richness of anurans. The final

models for each response variables were analyzed including but those with only significant main factors effect for anurans. Duncan multiple range test (DMRT) and Least square differences (LSD) were carried out to compare means of independent variables with significant variations at $p < 0.05$

RESULTS AND DISCUSSION

Anurans Species Richness

The species richness in the primary forest is significantly ($p < 0.05$) different from slash-and-burn cultivation (Fig. 2). Frog population increases steadily during natural succession, attaining similar characteristics to those from mature forest after regeneration following slash-and-burn cultivation (Pawar et al., 2004). Meanwhile, when the habitat starts to stabilize its condition (that is conducive to anurans), the frog species starts to increase dramatically.




Figure 2. Number of species of anurans observed in the different sites per collection period

Meanwhile, the total number of species in the primary forest, secondary forest, and slash-and-burn cultivation were $S=7$, $S=4$, and $S=0$, respectively. The result is consistent with the study of Mallari

et al. (2013) in Silago forest where they observed a maximum number of anurans species of $S=4$. Mallari et al. (2013) reported that the number of species is correlated to the degree of disturbance. Furthermore, the density of frogs observed was eight (8) frogs per 100 m² in the primary forest while four (4) frogs per 100 m² in the secondary forest and zero (0) frog in the slash-and-burn cultivation. This supports the study of Sluys et al. (2007) that the density observed in the rainforest in Brazil is four (4) frogs per 100 m². Moreover, it was observed that there were four (4) species of frogs (i.e., *Platymantis corrugatus*, *Occidozygalaavis*, *Platymantis* sp., and *Staurois*) found in primary forest that were not observed in the secondary forest and slash-and-burn cultivation while *Hylarana erythraea* (an introduced and invasive species) was observed in the secondary forest and was not observed in the other two sites.

Habitat Conditions that Influence Anurans Species Richness

Three environmental factors were considered in this study to determine the habitat condition of each site that influenced anurans species richness. These are microclimatic factors including temperature, relative humidity and rainfall (Cueto and Casenave 1999; Cushman, 2006), microhabitat structure including litter depth and quantity and quality of coarse woody debris

(deMaynadier and Hunter, 1995), and vegetation structure with the following layers, herbaceous layer, understorey and climax or emergent (Cramer and Willig 2002; Tews et al. 2004; Gingold et al. 2010; MacClain and Barry 2010).

Microclimatic Factors

In this study, three atmospheric factors (i.e., temperature, relative humidity, and precipitation) were measured that considerably affected the diversity of anurans in each study site. The results showed that temperature (during second and third collection) were significantly ($p < 0.01$) different in the three sites (Table 1). This is probably because the temperature in the primary and secondary forest sites is constantly stabilized by its multi-strata vegetation structure (Table 3). The result agrees with the findings of Markussen, (2005) where temperature in secondary forest is 2°C higher compared to primary forest. Hence, the result of this study supports the findings of Bickford, (2005) that anurans tend to seek cooler microhabitat with lower temperature. However, during the first collection, secondary forest and slash-and-burn cultivation had no significant difference. This is because during the measurement the sun was covered with clouds which prevented the radiation from penetrating to the land surface thereby stabilizing the temperature in both sites.

Table 1. Microclimatic factors measured at the different study sites per collection period

Climatic factors	Primary forest	Secondary forest	Slash and burn cultivation
1. Temperature (°C)			
Sampling 1	29.85±0.11 ^a	30.09±0.11 ^b	31.71±0.11 ^b
Sampling 2	25.27±0.03 ^a	25.87±0.03 ^b	27.17±0.03 ^c
Sampling 3	24.63±0.04 ^a	24.86±0.04 ^b	26.15±0.04 ^c
2. Relative Humidity (%)			
Sampling 1	83.74±0.71 ^a	80.69±0.71 ^b	67.81±0.71 ^c
Sampling 2	96.52±0.41 ^a	96.05±0.41 ^b	93.33±0.41 ^c
Sampling 3	96.09±0.16 ^a	95.52±0.16 ^b	93.84±0.16 ^c
3. Rainfall (cm week ⁻¹)			
Sampling 1	0.18±0.02 ^a	0.24±0.02 ^b	0.38±0.02 ^c
Sampling 2	0.89±0.01 ^a	0.92±0.01 ^b	1.23±0.01 ^b
Sampling 3	0.38±0.01 ^a	0.53±0.01 ^b	0.72±0.01 ^c

Note: least square means in each row with different letter superscript (a-c) are significantly different at p<0.05

The results for relative humidity showed that all sites were significantly ($p \leq .01$) different (Table 1). Reason might be due to vegetation structure which is obviously different in all study plots. Moreover, secondary forest has no dominant layer which might affect the retention of water vapor in the area. According to Bernatzky (1978), the stomata present in the leaves prevents the radiation to penetrate by absorbing and converting the heat energy into oxygen and glucose, thereby, minimizing the intensity of radiation which evaporates the water vapor rapidly resulting to higher relative humidity and lower temperature.

On the other hand, the rainfall during the collection period were significantly ($p \leq 0.01$) different in the three sites (Table 1). According

to Chazdon (2003), mean annual rainfall differs between primary and secondary forest. The high multi strata vegetation cover in the primary forest can trap more rain drops compared to vegetation layer with fewer strata in the secondary forest (Table 3). However, in the case of second collection, there was no significant difference between secondary forest and slash and burn. This could probably be due to the microclimate condition where some areas experience rain while some parts have none which is usually observed in tropical evergreen rainforest like in the study site. Also, the through fall or stem flow of the rain (which were not determine in the study) might contribute to the results.

Microhabitat Structure

The results showed that litter

depth was significantly different ($p < 0.01$) in each site (Table 2). Hence, the result implies that litter depth affects anurans richness (Fig.2). Several species of forest anurans are correlated with the quantity and quality of woody debris, litter depth, and over-storey canopy closure (deMaynadier and Hunter, 1995). According to Fauth et al. (1989), litter depth may

enhance diversity by providing a wider range of microhabitats and refuges against predators, as well as greater densities of arthropod preys. Moreover, for terrestrial breeding frogs, the leaf layer may provide adequate conditions for egg laying. This may facilitate the coexistence of more individuals and more species in the leaf litter (Sluys et al., 2007).

Table 2. Microhabitat structure measured at the different study sites

Microhabitat structure	Primary forest	Secondary forest	Slash and burn cultivation
Litter Depth (cm)	6.04±0.52 ^a	1.87±0.52 ^b	0.00±0.52 ^c
Quantity CWD	2.25±1.25 ^a	0.75±1.25 ^a	7.25±1.25 ^b
Quality CWD	2.75±0.56 ^a	1.63±0.56 ^b	1.00±0.56 ^c

Note: least square means in each row with different letter superscript (a-c) are significantly different at $p < 0.05$

Table 2 also shows that primary and secondary forest were significantly ($p < 0.01$) different with slash-and-burn cultivation in terms of the quantity of coarse woody debris. However, the high number of CWD in the slash-and-burn cultivation was due to the conversion of forest to agriculture where cutting and burning of trees were usually undertaken. Hence, the site was dominated by tree stumps and partially burned logs. The result of quality of CWD showed significant ($p < 0.01$) difference in all sites with highest observed value in the primary forest. This could probably because primary forest has higher decomposition rate. Therefore, the higher the quality of CWD, the greater the probability that anurans

use this as their habitat (Woodwall and Willams, 2005) which is supported by the data in Figure 2.

Vegetation Structure

Table 3 shows a significant difference ($p < 0.01$) in the understory layer for all site. The high number of understory in the secondary forest was because it is on its early successional stage where most small pioneer trees and shrubs dominate the understory layer. Likewise, herbaceous layer had significant difference ($p < 0.01$) in all sites. The high number of herbaceous layer observed in the secondary forest was due to the community of grass and some ferns dominating the site while in the primary forest was because of wildlings from the trees.

Table 3. Vegetation structure measured at the different study sites

Vegetation strata	Primary forest	Secondary forest	Slash and burn cultivation
Emergent	4.25±8.54 ^a	0.00±8.54 ^b	0.00±8.54 ^b
Understorey	12.50±1.67 ^a	19.25±1.67 ^b	0.00±1.67 ^c
Herbaceous Layer	50.00±0.28 ^a	106.50±0.28 ^b	0.00±0.28 ^c

Note: least square means in each row with different letter superscript (a-c) are significantly different at $p < 0.05$

According to Feder (1983), anuran species maintain close contact with forest floor substrates and habitat selection are also affected by any anthropogenic activities. The pattern of increase in habitat heterogeneity on the structure and composition of vegetation becomes complex if the niche diversity and species diversity increases (Cramer and Willig, 2002; Tews et al., 2004; Gingold et al., 2010; McClain and Barry, 2010). This implies that clearing the vegetation would disturb the area; thus, greatly affecting the habitat condition and eventually the species richness of the frogs.

CONCLUSION

The results of the study affirm that anthropogenic disturbance alters anurans habitat condition, thereby, affecting its species richness and population. The study was not able to find specific species of anurans that will be used as indicator to ecological disturbance. However, it is worth mentioning that there was a considerable reduction ($p > 0.05$) of anurans species in the primary forest, secondary forest, and slash-and-burn cultivation with values of seven (7), four (4) and zero (0)

respectively. Moreover, the study suggests that microclimatic parameters, microhabitat, and vegetation structures appear to be the factors influencing habitat condition of anurans which significantly ($p > 0.05$) affect anurans species richness.

RECOMMENDATION

The applicability of different methods on the sampling of anurans varies depending on the aim of the study. Therefore, you need to assess first the main objective of the study before selecting a specific method. Also, the duration of the collection period needs to be increased to also increase data variability and thereby could draw a conclusive statement. Furthermore, a study on the insect composition should also be conducted in order to support the reason of the decrease of anurans was due to the disturbance and not because of the unavailability of food.

It is also recommended that proper management of habitat needs to be implemented in the area for the conservation of anurans. Likewise, studying the requirement of habitat preference of a specific frog is essential for the management of their habitat.

REFERENCES:

- ALCALA, A.C. and W.C. BROWN. 1996. Philippine Amphibians: An Illustrated Field Guide. Bookmark Inc., Makati City, Philippines.
- ALCALA, A.C. 1986. Guide to Philippine Flora and Fauna: Vol X, Amphibians and Reptiles. Natural Resource Management Centre, Ministry of Natural Resources and the University of the Philippines, Manila, Philippines.
- ALVAREZ, D. and A.G. NICIEZA. 2002. Effects of Temperature and Food Quality on Larval Growth and Metamorphosis. *Fuente Ecol* 16:640-648
- BICKFORD, D. 2005. Long-term frog monitoring with local people in Papua New Guinea and the 1997-98 El Niño Southern Oscillation Event. In: Donnelly M, White M, Crother B, Wake C (eds) *Ecology and evolution in the tropics—a herpetological perspective*. University of Chicago Press, Chicago
- CENIZA, M. J., BANDE M. J., FERNANDO, E. S., LABASTILLA, P. K., COME, R. S., OMEGA, R. G., ALESNA, W. T., 2011. Community-based forest restoration and biodiversity protection and management of lowland dipterocarp forests in Silago,
- CHAZDON, R. L., 2003. Tropical forest recovery: legacies of human impact and natural disturbances. Vol. 6/1, 2, pp. 51-71. Urban & Fischer Verlag
- CRAMER, M.J., WILLIG, M.R. 2002: Habitat heterogeneity, habitat associations, and rodent species diversity in a sand-shinnery-oak landscape. *Journal of Mammalogy*, 743-753.
- CUETO, V.R., DE CASENAVE, J.L. 1999: Determinants of bird species richness: Role of climate and vegetation structure at a regional scale. *Journal of Biogeography*, 26(3), 487-492. Published by Blackwell Publishing.
- CUSHMAN, S.A. 2006. Effects of habitat loss and fragmentation on amphibians: A review and prospectus. *Biological Conservation*. 128:231-240
- DEY, M. 2010. A Study on Habitat Selection and Probable Anthropogenic Threats of Anurans in Barak Valley, Northeast India.
- DEMAYNADIER, P.G. and M.L. JR., HUNTER. 1995. The Relationship between Forest Management and Amphibian Ecology: A Review of the North American Literature. *Environmental Reviews* 3(3&4):230-261.
- DIESMOS, A., A. ALCALA, R. BROWN, L. AFUANG, C. DOLINO, G. GEE, K. HAMPSON, M.L. DIESMOS, A. MALLARI, P. ONG, L. PAGUNTALAN, M. PEDREGOSA, D. UBALDO, B. GUTIERREZ. 2004. *Platymantis corrugatus*. In: IUCN 2012
- FAUTH J. E., CAROTHER B. I. & SLOWINSKI J. B. (1989) Elevational patterns of species richness, evenness and abundance of the Costa Rican leaf-litter herpetofauna. *Biotropica* 21, 178-85.
- FEDER, M. E. 1983. Integrating the Ecology and Physiology of Plethodontid Salamanders. *Herpetologica* 39:291-310.
- GINGOLD, R., MUNDO-OCAMPO, M., HOLOVACHOV, O., ROCHA-OLIVARES, A. 2010: The role of habitat heterogeneity in structuring the community of intertidal free-living marine nematodes. *Marine Biology: International Journal on Life in Oceans and Coastal Waters*, 157(8), 1741-1753. doi: 10.1007/s00227-010-1447-z. Retrieved from: <http://escholarship.org/uc/item/6qp3406w>
- HEYING. 2004. Introduction to Amphibians. *Heyinglect 4a-FTTS Fall 2004 (H5)*

- INTERNATIONAL UNION FOR CONSERVATION OF NATURE (IUCN), 2015. The IUCN Red List of Threatened Species version 2015.
- KUMMER, D.M. 1992. Deforestation in the Postwar Philippines. Chicago: University of Chicago Press, 1992(a): 1-177.
- MACCLAIN, C.R., BARRY, J.P. 2010: Habitat heterogeneity, disturbance, and productivity work in concert to regulate biodiversity in deep submarine canyons. Ecology, Ecological Society of America, 964-976.
- MARKUSSEN, M. 2005. Valuation and conservation of biodiversity; Interdisciplinary perspective on convention in biological diversity. ISBN 3-540-24022-5 Springer Berlin Heidelberg New York.
- MALLARI, N. A., DIEMOS, A., SUPSOP, C., and ALBAN, J. D., 2013. Biodiversity Baseline Assessment in the REDDplus Pilot Area on Leyte Island as an Input for the Elaboration of a MRV system for REDD-plus including biodiversity co-benefits. A technical report published by; Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ) GmbH
- MALLARI, N.A.D. 2009: Maximising the value of ecological and socio-economic data in support of conservation planning for key understorey bird species in Palawan, Philippines. Ph. D. Dissertation. Department of Environmental and Geographical Sciences. Manchester Metropolitan University. Unpublished
- MARGRAF, J. and P. P. MILAN 2004. Rainforestation Farming. A Farmer's Guide to Biodiversity Management for the Philippines. 1st ed., Leyte State University, Baybay, Leyte, Philippines.
- MYERS, N., R.A. MITTERMEIER, C.G. MITTERMEIER. 2000 Biodiversity hotspots for conservation priorities. Nature 403:853-858
- PAWAR, S. S., RAWAT, G. S., CHOUDHURY, B. C. 2004. Recovery of frog and lizard communities following primary habitat alteration in Mizoram, Northeast India.
- SLUYS, M. V., VRCIBRADIC, D., ALVES, M. A., BERGALLO, H. G. and. ROCHA, C. D. 2007. Ecological parameters of the leaf-litter frog community of an Atlantic Rainforest area at Ilha Grande, Rio de Janeiro state, Brazil. Austral Ecology (2007) 32, 254-260
- SHULTE, A. 2002. Rainforestation Farming: Option for Rural Development and Biodiversity Conversion in the Humid Tropics of Southeast Asia: A Review on Major Issues on Community-Based Rehabilitation Silvicultural and Guide to Economic Tree Species in the Visayas/Philippines.
- SVENSSON, R.J. 2010. Ecological disturbances: The Good, the Bad and the Ugly. Geson Hylte Tryck, Göteborg, Sweden. ISBN: 978-91-628-8200-6.
- TEWS, J., BROSE, U., GRIMM, V., TIELBÖRGER, K., WICHMANN, M.C., SCHWAGER, M., JELTSCH, F. 2004: Animal species diversity driven by habitat heterogeneity/diversity: The importance of keystone structures. Journal of Biogeography, 31(1), 79-92. doi: 10.1046/j.0305-0270.2003.00994.x
- WARGUEZ, D. A., MONDEJAR, E. P. and DEMAYO, C. G. 2013. Frogs and their Microhabitat Preferences in the Agricultural and Secondary Forest areas in the Vicinity of Mt. Kalatungan Mountain, Bukidnon, Philippines. International Research Journal of Biological Science. ISSN 2278-3202.
- WILLIAMS, L. A. 2004 Amphibian population and community characteristics habitat relationship, and first-year response to clearcutting in a central Appalachian Industrial
- WOODWALL C. and WILLIAMS M.S. 2005. Sampling protocol, estimation and analysis procedure for the down woody materials indicator of the FIA program. General Technical Report NC-256. St. Paul, MN: US, Department of Agriculture, Forest Service, North Central Research Station, 47 pp.